


REVIEW, pages 330–335

4.1

1. a) Use a table of values to graph $y = 2x^2 + 6x - 8$.

x	-5	-4	-3	-2	-1	0	1	2
y	12	0	-8	-12	-12	-8	0	12



b) Determine:

- i) the intercepts
- ii) the coordinates of the vertex
- iii) the equation of the axis of symmetry
- iv) the domain of the function
- v) the range of the function

Give the values to the nearest tenth where necessary.

i) **x-intercepts:** $-4, 1$

y-intercept: -8

ii) **From the graph, the axis of symmetry is midway between $x = -1$ and $x = -2$.**

So, the equation of the axis of symmetry is $x = -1.5$.

When $x = -1.5$, $y = 2(-1.5)^2 + 6(-1.5) - 8$, or -12.5

The coordinates of the vertex are: $(-1.5, -12.5)$

iii) **axis of symmetry:** $x = -1.5$

iv) **domain:** $x \in \mathbb{R}$

v) **range:** $y \geq -12.5, y \in \mathbb{R}$

2. Which of these tables of values represents a quadratic function?

Justify your response.

a)

x	0	1	2	3
y	3	-3	-13	-27

The x-coordinates increase by 1 each time.

First differences:

$$-3 - 3 = -6$$

$$-13 - (-3) = -10$$

$$-27 - (-13) = -14$$

The first differences decrease by 4 each time. So, the function is quadratic.

b)

x	0	1	2	3
y	1	3	5	7

The x-coordinates increase by 1 each time.

First differences:

$$3 - 1 = 2$$

$$5 - 3 = 2$$

$$7 - 5 = 2$$

The first differences are constant. So, the function is linear.

4.2

3. Use graphing technology to approximate the solution of the equation below. Write the roots to 3 decimal places.

$$3x^2 + 6x - 70 = 0$$

Graph $y = 3x^2 + 6x - 70$. Use the CALC feature to display $X = -5.932883$ and $X = 3.9328829$. The roots are approximately $x = -5.933$ and $x = 3.933$.

4.3

4. For each pair of quadratic functions, describe how their graphs are related.

a) $y = (x - 3)^2$; $y = (x + 2)^2$

Compare the equations with $y = (x - p)^2$.

$y = (x - 3)^2$: Its graph is the graph of $y = x^2$ translated 3 units to the right.

$y = (x + 2)^2$: Its graph is the graph of $y = x^2$ translated 2 units to the left.

So, the graph of $y = (x - 3)^2$ is translated 5 units left to get the graph of $y = (x + 2)^2$.

b) $y = x^2 + 5$; $y = x^2 - 1$

Compare the equations with $y = x^2 + q$.

$y = x^2 + 5$: Its graph is the graph of $y = x^2$ translated 5 units up.

$y = x^2 - 1$: Its graph is the graph of $y = x^2$ translated 1 unit down.

So, the graph of $y = x^2 + 5$ is translated 6 units down to get the graph of $y = x^2 - 1$.

c) $y = -\frac{1}{2}x^2$; $y = \frac{1}{2}x^2$

Compare the equations with $y = ax^2$.

$y = -\frac{1}{2}x^2$: Its graph is the graph of $y = x^2$ compressed by a vertical factor of $\frac{1}{2}$, then reflected in the x -axis.

$y = \frac{1}{2}x^2$: Its graph is the graph of $y = x^2$ compressed by a vertical factor of $\frac{1}{2}$.

So, the graph of $y = -\frac{1}{2}x^2$ is reflected in the x -axis to get the graph of $y = \frac{1}{2}x^2$.

4.4

5. For this quadratic function: $y = \frac{1}{2}(x - 4)^2 - 2$

- a) Identify the coordinates of the vertex, the domain, the range, the direction of opening, the equation of the axis of symmetry, and the intercepts.

a is positive, so the graph opens up.

$p = 4$ and $q = -2$, so the coordinates of the vertex are: $(4, -2)$

The equation of the axis of symmetry is $x = p$; that is, $x = 4$.

To determine the y -intercept, substitute $x = 0$:

$$y = \frac{1}{2}(0 - 4)^2 - 2$$

$$y = 6$$

The y -intercept is 6.

To determine the x -intercepts, substitute $y = 0$:

$$0 = \frac{1}{2}(x - 4)^2 - 2$$

$$0 = \frac{x^2}{2} - 4x + 6$$

$$0 = x^2 - 8x + 12$$

$$0 = (x - 6)(x - 2)$$

$$x = 6 \text{ or } x = 2$$

The x -intercepts are 2 and 6.

The domain is: $x \in \mathbb{R}$


The graph opens up, so the vertex is a minimum point with y -coordinate -2 . The range is: $y \geq -2, y \in \mathbb{R}$

- b) Sketch a graph.

The graph is congruent to the graph of $y = \frac{1}{2}x^2$.

On a grid, mark a point at the vertex $(4, -2)$. Use the step pattern.

Multiply each vertical step by $\frac{1}{2}$.



6. Determine an equation of the quadratic function for each set of data given.

a) The coordinates of the vertex are $V(4, 12)$ and the graph passes through $A(7, 6)$.

An equation has the form $y = a(x - p)^2 + q$.

The vertex is at $V(4, 12)$, so $p = 4$ and $q = 12$.

The equation becomes $y = a(x - 4)^2 + 12$.

Substitute the given coordinates for point A: $x = 7, y = 6$

$$6 = a(7 - 4)^2 + 12$$

$$6 = 9a + 12$$

$$-6 = 9a$$

$$a = -\frac{2}{3}$$

So, the equation of the function is:

$$y = -\frac{2}{3}(x - 4)^2 + 12$$

b) The graph passes through $B(2, -5)$ and has x -intercepts -3 and 4 .

Use $y = a(x - x_1)(x - x_2)$ Substitute: $x_1 = -3$ and $x_2 = 4$

$y = a(x + 3)(x - 4)$ Substitute for $B(2, -5)$.

$$-5 = a(2 + 3)(2 - 4)$$

$$-5 = -10a$$

$$a = 0.5$$

In factored form, the equation is: $y = 0.5(x + 3)(x - 4)$

4.5

7. Write this equation in standard form.

$$y = -3x^2 + 24x - 45$$

$$y = -3(x^2 - 8x) - 45$$

$$\text{Add and subtract: } \left(\frac{-8}{2}\right)^2 = 16$$

$$\begin{aligned} y &= -3(x^2 - 8x) - 45 \\ &= -3(x^2 - 8x + 16 - 16) - 45 \\ &= -3(x^2 - 8x + 16) + 48 - 45 \\ &= -3(x - 4)^2 + 3 \end{aligned}$$

4.6

8. Determine the intercepts, the equation of the axis of symmetry, and the coordinates of the vertex of the graph of each quadratic function, then sketch the graph.

a) $y = 2x^2 + 2x - 24$

The y -intercept is -24 .

Factor the equation.

$$\begin{aligned} y &= 2x^2 + 2x - 24 \\ &= 2(x^2 + x - 12) \\ &= 2(x + 4)(x - 3) \end{aligned}$$

The x -intercepts are: -4 and 3

The mean of the intercepts is:


$$\frac{-4 + 3}{2} = -0.5$$

So, the equation of the axis of symmetry is: $x = -0.5$

Substitute $x = -0.5$ in

$$\begin{aligned} y &= 2x^2 + 2x - 24 \\ &= 2(-0.5)^2 + 2(-0.5) - 24 \\ &= -24.5 \end{aligned}$$

The coordinates of the vertex are: $(-0.5, -24.5)$



b) $y = -\frac{1}{2}x^2 - x + 4$

The y -intercept is 4 .

Factor the equation.

$$\begin{aligned} y &= -\frac{1}{2}x^2 - x + 4 \\ &= -\frac{1}{2}(x^2 + 2x - 8) \\ &= -\frac{1}{2}(x + 4)(x - 2) \end{aligned}$$

The x -intercepts are: -4 and 2

The mean of the intercepts is:


$$\frac{-4 + 2}{2} = -1$$

So, the equation of the axis of symmetry is: $x = -1$

Substitute $x = -1$ in

$$\begin{aligned} y &= -\frac{1}{2}x^2 - x + 4 \\ &= -\frac{1}{2}(-1)^2 + 1 + 4 \\ &= 4.5 \end{aligned}$$

The coordinates of the vertex are: $(-1, 4.5)$



4.7

9. Select Audio Company sells an MP3 player for \$75. At that price, the company sells approximately 1000 players per week. The company predicts that for every \$5 increase in price, it will sell 50 fewer MP3 players. Which price for an MP3 player will maximize the revenue?

Let x represent the number of \$5 increases in the price of an MP3 player.

When the cost is \$75, 1000 are sold for a revenue of:

$$\$75(1000) = \$75\,000$$

When the cost is $\$(75 + 5x)$, $(1000 - 50x)$ are sold for a revenue of $\$(75 + 5x)(1000 - 50x)$.

Let the revenue be R dollars.

An equation is: $R = (75 + 5x)(1000 - 50x)$

Use a graphing calculator to graph the equation.

From the graph, the maximum revenue is about \$76 562.50 when the number of \$5 increases is 2.5.

The number of increases is a whole number, so round 2.5 to 2 or to 3.

Two increases of \$5 mean that the MP3 player will now cost:

$$2(\$5) + \$75 = \$85$$

Three increases of \$5 mean that the MP3 player will now cost:

$$3(\$5) + \$75 = \$90$$

To maximize the revenue, the MP3 player should sell for \$85 or \$90.